

$B \rightarrow X_s \tau^+ \tau^-$ in the flipped SU(5) model

Chao-Shang Huang and Qi-Shu Yan
 Institute of Theoretical Physics, Academia Sinica.
 P.O. Box 2735, Beijing 100080, P.R. China

Abstract

We perform a detailed study of the inclusive branching ratio and the forward-backward asymmetry of the rare B decay process $B \rightarrow X_s \tau^+ \tau^-$ in the flipped SU(5) model, a predictive model which has only two free parameters, plus the sign of the higgs mixing mass μ , to describe mass spectrum and mixings of about 30 sparticles. In contrast to other works dealing with such a topic, our study has included the contributions of the neutral higgs bosons through penguin diagrams to this process under the context of supersymmetry. In some regions of the parameter space, due to the substantial enhancement effects coming from supersymmetry, these contributions become quite significant and could make the flavor change process $B \rightarrow X_s \tau^+ \tau^-$ a microscope to probe the regions of large $\tan\beta$ and a window to gain an insight into new physics beyond the standard model(SM).

1 Introduction

The standard model(SM) is a successful effective low energy theory at energy scales up to 100 GeV; its predictions are in agreement with almost all experimental tests. Nevertheless, there exist many theoretical problems waiting for explanations. For example, where do Yukawa couplings originate from? Why are there three, not other numbers of, generations? Supersymmetry is an excellent candidate which offers a scheme to embed the SM in a more fundamental theory with which these theoretical problems can be hopefully explained. Supersymmetry has many good features. It ingeniously tackles the abominable gauge hierarchy problem from which ordinary grand unification theories suffer [1], provides a mechanism that supergravity models always share [2] which dynamically breaks the electroweak symmetry via radiative corrections and avoids the arbitrary procedure to break $SU(2) \times U(1)$ group by hand, furnishes an elegant framework to unify gravitational, strong and electroweak interactions [3], naturally proposes the lightest supersymmetric particle(LSP) as a candidate of the dark matter in astrophysics and cosmology [4], and even can explain all parameters and properties of SM in principles with only one input, as many superstring models have demonstrated [5]. For all these compelling features, supersymmetry faces its biggest problem: so far there are no decisive evidences affirming its existence.

At present, in spite of lacking definite supports from experiments, supersymmetry could be understood in an experimental perspective. Upon various possible new physics beyond SM, latest experimental results impose very severe constraints; supersymmetry must confront these tests, as summarized in [6]. About SUSY we know: 1)charged SUSY particles are heavier than 65 GeV, as given by the experiments at LEP [7]; 2)the gluino mass is larger than 100 GeV, a bound given by Fermilab Tevatron collider [8]; 3)all sneutrino masses are no less than 41 GeV [9]; 4)there should exist no seeds to break charge and color symmetry [10]; 5)the lightest supersymmetric particle is neutral; 6)the width of the process $Z \rightarrow \chi\chi$ is less than 8.4 MeV, and branching ratios of $Z \rightarrow \chi\chi'$ and $Z \rightarrow \chi'\chi'$ are less than 2×10^{-5} ; 7)the sparticles masses should not be much larger than 1TeV [11], otherwise the higgs bosons loop corrections would be too large and gauge hierarchy problem arise again; 8)the most stringent constraint is brought in by the branching ratio of the rare decay $b \rightarrow s\gamma$, the value is determined to be $(2.32 \pm 0.57 \pm 0.35 \times 10^{-4})$ with the upper bound 4.2×10^{-4} and lower bound 1×10^{-4} at CLEO [12].

There are many supersymmetric models to be tested. For experimentalists, to test a generic supersymmetric model, the minimal supersymmetrically extended standard model without constraints on soft breaking parameters for instance, makes no much sense, because so many possible explanations for a test can be offered by the model that no significant predictions can be made for other tests. In contrast to this kind of models, a well theoretically motivated model is highly predictive, therefore easier to be proven true or false, and so is favored by experimentalists. Among various SUSY models, the flipped SU(5) model [13, 14], one of a class of string-inspired supergravity models, is such a predictive model. In the model the proton decay and cosmological constraints are satisfied automatically and there are only two free parameters (plus the sign of the higgs

mixing mass μ): the gaugino mass evaluated at the unification scale $m_{1/2}$ and the ratio of the higgs vacuum expectation values $\tan\beta$. Constraints from phenomenology listed above greatly cut down the parameter spaces permitted by the model. In this note we shall study $B \rightarrow X_s \tau^+ \tau^-$ in the model.

As analyzed in literatures [15, 16], $b \rightarrow s\gamma$ puts a stringent constraint on various possible new physics beyond SM. For example, in the two-higgs-doublet model, it requires a larger charged higgs mass. In SUSY, this restrictive condition can be relaxed or be grimmer due to the either destructive or constructive interferences between the contributions of charginos and those of $H^\pm$ and $W^\pm$ [14, 15, 16]. In the flipped SU(5) model, for the case $\mu > 0$, it can drastically reduce the parameter space, especially the regions of permitted $\tan\beta$, while for the case $\mu < 0$, it imposes no such a stringent constraint upon $\tan\beta$ [14].

Compared with the process $b \rightarrow s\gamma$ whose magnitude of branching ratio(BR) is 10^{-4} , observing another rare B decay process $b \rightarrow sl^+l^-$ (its BR is 10^{-6} [17]) is not available at present. But with the advent of the construction of B factories, this process is also quite promising and, because of its sensitivity to new physics, has been proposed as one powerful piece of the arsenal to discern supersymmetry from SM [17, 18]. It was found in the minimal supergravity model that this process is strongly correlated with $b \rightarrow s\gamma$ and in some regions of the parameter space the branching ratio of it can be enhanced by about 50% compared with SM [6]. All the considerations above have ignored the contributions from exchanging neutral higgs bosons(NHB). In SM, they can be safely neglected owing to the smallness of m_l/m_w ($l=e, \mu, \tau$). But it is unjustifiable to omit them when beyond SM and for the process $b \rightarrow s\tau^+\tau^-$. As pointed out in [19], owing to the enhancement effects of the large $\tan\beta$ in two-higgs-doublet model, they become quite sizable and thus greatly influence the invariant mass distribution and forward-backward asymmetry of $b \rightarrow s\tau^+\tau^-$. Under the context of SUSY, as we shall show below, a more significant enhancement effect of large $\tan\beta$ coming from chargino-stop loops can even make them dominant contributions so that in some regions of the parameter space the branching ratio of $b \rightarrow s\tau^+\tau^-$ can be enhanced by 200% in both scenarios when compared with SM. Therefore, it is possible that the first distinct signs of supersymmetry could come from deviations in $b \rightarrow sl^+l^-$, especially, for $l=\tau$.

2 A brief description of the model

The detailed description of the string inspired flipped SU(5) model can be found in [13]. Here we only discuss two aspects of it which are relevant to the study of $B \rightarrow X_s \tau^+ \tau^-$ in this paper. One is about the unification of gauge couplings at string determined scale, the other the number of the free parameters.

It is well known that the scale of gauge coupling unification derived from string theory is about 7×10^{17} GeV [20, 21], much larger than the experimentally determined value 2×10^{16} GeV, when assuming the particle content of the minimal supersymmetric standard model. Many approaches have been proposed to reconcile the discrepancy between these

two scales [22]. In the flipped $SU(5)$ grand unified theory, this problem was solved by introducing one pairs of gap particles(GPs) which can be economically embedded into the representations of $SU(5) \times U(1)$ [13]. They decouple from interactions later than heavier particles and bridge the gap between these two scales.

Universal soft supersymmetric terms can be derived from local and global supersymmetry breaking. Four free parameters, all are evaluated at unified scale, are introduced here: $m_{1/2}$, the mass of gauginos; A , the trilinear couplings; B , the bilinear couplings; m_0 , the universal masses for all scalars. String-inspired relations lead to two supersymmetry breaking scenarios: 1)the moduli scenario which requires $m_0=A=0$ [2]; 2)the dilaton scenario which requires $m_0=m_{1/2}/\sqrt{3}$, $A=-m_{1/2}$ [23]. Thus only B and $m_{1/2}$ are the free soft breaking parameters which will be further constrained.

Electroweak symmetry is dynamically broken at low energy due to the fact that the mass of one of the two higgs fields becomes negative because of the effect of the large value of the top's Yukawa coupling. This is known as the radiative electroweak symmetry breaking mechanism. Minimal conditions of the scalar potential give

$$\begin{aligned}\sin 2\beta &= \frac{-2B\mu}{m_{H_1}^2 + m_{H_2}^2 + 2\mu^2} \\ \tan^2\beta &= \frac{m_{H_1}^2 + \mu^2 + m_z^2/2}{m_{H_2}^2 + \mu^2 + m_z^2/2}\end{aligned}\tag{1}$$

which go a further step to reduce the number of free parameters, for B and μ^2 can be solved out from these two equations while leaving the sign of the higgs mixing parameter μ undetermined. Thus only two free parameters ($m_{1/2}$ and $\tan\beta$) survive to completely describe mass spectra and mixings of about 30 sparticles.

We use eq.(1) and the renormalization group equations(RGEs) of masses and couplings [24] to calculate mass spectra and mixings. In order to simplify the computational procedure, we neglect the Yukawa couplings of the first two generations and the effects of GPs. A detailed analysis which includes these effects will be published elsewhere. With the constraints from phenomenology we take the range of $\tan\beta$ as from 2 to 40, and $m_{1/2}$ from 100 GeV to 400 GeV. The constraint from $b \rightarrow s\gamma$ will be taken into account later on in our analysis.

3 The formula for $B \rightarrow X_s \tau^+ \tau^-$

Inclusive decay rates of heavy hadrons can be calculated in heavy quark effective theory and it has been shown that the leading terms in $1/m_Q$ expansions turns to be the decay of a free (heavy) quark and corrections stem from the order $1/m_Q^2$ [25]. In what follows we shall calculate the leading term.

Under the context of supersymmetry, there are total five classes of loops contributing to the flavor changing process $b \rightarrow s \tau^+ \tau^-$ as well as to $b \rightarrow s\gamma$ by exchanging: W boson and u-type quarks, charged higgs boson and u-type quarks, charginos and u-type squarks, neutralinos and d-type squarks, gluinos and d-type squarks. Since the flavor mixings

between the third and the other two generations are small, contributions from the last two cases are negligible so that we only consider contributions from the first three classes. The additional Feynman diagrams are shown in fig.1.

With Feynman rules at electroweak scale, as given in [26], effective weak Hamiltonian can be obtained

$$H_{eff} = \frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \left(\sum_{i=1}^{10} C_i(\mu) O_i(\mu) + \sum_{i=1}^{10} C_{Q_i}(\mu) Q_i(\mu) \right) \quad (2)$$

where O_i and Q_i are given in [27] and [19] respectively. The coefficients $C_i(m_W)$ can be found in [17]. We calculate the coefficients $C_{Q_i}(m_W)$ in SUSY models and the results are:

$$\begin{aligned} C_{Q1}(m_W) &= \frac{m_b m_\tau}{4m_{h_0}^2 \sin^2 \theta_W} \text{tg}^2 \beta \{ (\sin^2 \alpha + h \cos^2 \alpha) \left[\frac{1}{x_{Wt}} (f_1(x_{Ht}) - f_1(x_{Wt})) \right. \right. \\ &+ \sqrt{2} \sum_{i=1}^2 \frac{m_{\chi_i}}{m_W \cos \beta} (-V_{i1} f_1(x_{\chi_i \tilde{q}}) + \sum_{k=1}^2 \Lambda(i, k) T_{k1} f_1(x_{\chi_i \tilde{t}_k})) \\ &+ (1 + \frac{m_{H^\pm}^2}{m_W^2}) f_2(x_{Ht}, x_{Wt})] - \frac{m_{h_0}^2}{m_W^2} f_2(x_{Ht}, x_{Wt}) \\ &+ 2 \sum_{ii'=1}^2 (B_1(i, i') \Gamma_1(i, i') + A_1(i, i') \Gamma_2(i, i')) \} \\ C_{Q2}(m_W) &= -\frac{m_b m_\tau}{4m_{A^0}^2 \sin^2 \theta_W} \text{tg}^2 \beta \{ \frac{1}{x_{Wt}} (f_1(x_{Ht}) - f_1(x_{Wt})) + 2f_2(x_{Ht}, x_{Wt}) \\ &+ \sqrt{2} \sum_{i=1}^2 \frac{m_{\chi_i}}{m_W \cos \beta} (-V_{i1} f_1(x_{\chi_i \tilde{q}}) + \sum_{k=1}^2 \Lambda(i, k) T_{k1} f_1(x_{\chi_i \tilde{t}_k})) \\ &+ 2 \sum_{ii'=1}^2 (-U_{i'2} V_{i1} \Gamma_1(i, i') + U_{i2}^* V_{i'1}^* \Gamma_2(i, i')) \} \end{aligned} \quad (3)$$

where

$$\begin{aligned} B_1(i, i') &= (-\frac{1}{2} U_{i'1} V_{i2} \sin 2\alpha (1 - h) + U_{i'2} V_{i1} (\sin^2 \alpha + h \cos^2 \alpha)) \\ A_1(i, i') &= (-\frac{1}{2} U_{i1}^* V_{i'2}^* \sin 2\alpha (1 - h) + U_{i2}^* V_{i'1}^* (\sin^2 \alpha + h \cos^2 \alpha)) \\ \Gamma_1(i, i') &= m_{\chi_i} m_{\chi_{i'}} U_{i2} (-\frac{1}{\tilde{m}^2} f_2(x_{\chi_i \tilde{q}}, x_{\chi_{i'} \tilde{q}}) V_{i'1} + \sum_{k=1}^2 \frac{1}{m_{\tilde{t}_k}^2} \Lambda(i', k) T_{k1} f_2(x_{\chi_i \tilde{t}_k}, x_{\chi_{i'} \tilde{t}_k})) \\ \Gamma_2(i, i') &= U_{i2} (-f_2(x_{\chi_i \tilde{q}}, x_{\chi_{i'} \tilde{q}}) V_{i'1} + \sum_{k=1}^2 \Lambda(i', k) T_{k1} f_2(x_{\chi_i \tilde{t}_k}, x_{\chi_{i'} \tilde{t}_k})) \\ \Lambda(i, k) &= V_{i1} T_{k1} - V_{i2} T_{k2} \frac{m_t}{\sqrt{2} m_W \sin \beta} \\ f_1(x_{ij}) &= 1 - \frac{x_{ij}}{x_{ij} - 1} \ln x_{ij} + \ln x_{Wj} \end{aligned}$$

$$\begin{aligned}
f_2(x, y) &= \frac{1}{x-y} \left(\frac{x}{x-1} \ln x - \frac{y}{y-1} \ln y \right) \\
x_{ij} &= m_i^2/m_j^2
\end{aligned} \tag{4}$$

with m_i being the mass of the particle i , and

$$\begin{aligned}
C_{Q3}(m_w) &= \frac{m_b e^2}{m_\tau g^2} \{C_{Q1}(m_w) + C_{Q2}(m_w)\} \\
C_{Q4}(m_w) &= \frac{m_b e^2}{m_\tau g^2} \{C_{Q1}(m_w) - C_{Q2}(m_w)\} \\
C_{Qi}(m_w) &= 0, (i = 5, \dots, 10)
\end{aligned} \tag{5}$$

In eqs.(3) and (4), U and V are matrices which diagonalize the mass matrix of charginos, T is the matrix reflecting the mixing of stops t_R and t_L , m_{χ_j} denote the chargino masses, $\tilde{m}$ is the average mass of u-type squarks $\tilde{q}$ of the first two generations, h is the square of the ratio of the mass of h^0 to the mass of H^0 and α is the mixing angle of neutral components of the two higgs doublets in the model. And in eq.(3) we have omitted less important terms because they are numerically negligible compared to those given in eq.(3) when $\tan\beta \geq 20$.

Considering QCD corrections and evolving these coefficients down to the scale we interest in, the effective Hamiltonian results in the following matrix elements for $B \rightarrow X_s \tau^+ \tau^-$

$$\begin{aligned}
M &= \frac{G_F \alpha}{\sqrt{2} \pi} V_{tb} V_{ts}^* [C_8^{eff} \bar{s}_L \gamma_\mu b_L \bar{\tau} \gamma^\mu \tau + C_9 \bar{s}_L \gamma_\mu b_L \bar{\tau} \gamma^\mu \gamma^5 \tau \\
&\quad + 2C_7 m_b \bar{s}_L i \sigma^{\mu\nu} \frac{q^\mu}{q^2} b_R \bar{\tau} \gamma^\nu \tau + C_{Q1} \bar{s}_L b_R \bar{\tau} \tau + C_{Q2} \bar{s}_L b_R \bar{\tau} \gamma^5 \tau]
\end{aligned} \tag{6}$$

here these coefficients are evaluated at $\mu=m_b$. C_8^{eff} is given as[27]:

$$C_8^{eff} = C_8 + \left\{ g\left(\frac{m_c}{m_b}, s\right) + \frac{3}{\alpha^2} k \sum_{V_i} \frac{\pi M_{V_i} \Gamma(V_i \rightarrow \tau^+ \tau^-)}{M_{V_i}^2 - q^2 - i M_{V_i} \Gamma_{V_i}} \right\} (3C_1 + C_2) \tag{7}$$

From eq.(6), by integrating the angle variable of the double differential distributions from 0 to π , the invariant dilepton mass distributions can be calculated and given below

$$\begin{aligned}
\frac{d\Gamma(B \rightarrow X_s \tau^+ \tau^-)}{ds} &= B(B \rightarrow X_c l \bar{\nu}) \frac{\alpha^2}{4\pi^2 f(m_c/m_b)} (1-s)^2 \left(1 - \frac{4t^2}{s}\right)^{1/2} \frac{|V_{tb} V_{ts}^*|^2}{|V_{cb}|^2} D(s) \\
D(s) &= |C_8^{eff}|^2 \left(1 + \frac{2t^2}{s}\right) (1+2s) + 4|C_7|^2 \left(1 + \frac{2t^2}{s}\right) \left(1 + \frac{2}{s}\right) \\
&\quad + |C_9|^2 \left[(1+2s) + \frac{2t^2}{s} (1-4s)\right] + 12\text{Re}(C_7 C_8^{eff*}) \left(1 + \frac{2t^2}{s}\right) \\
&\quad + \frac{3}{2} |C_{Q1}|^2 (s - 4t^2) + \frac{3}{2} |C_{Q2}|^2 s + 6\text{Re}(C_9 C_{Q2}^*) t
\end{aligned} \tag{8}$$

where $s=q^2/m_b^2$, $t=m_\tau/m_b$, $B(B \rightarrow X_c l \bar{\nu})$ is the branching ratio which takes as 0.11, f is the phase-space factor and $f(x)=1-8x^2+8x^6-x^8-24x^4 \ln x$. The forward-backward asymmetry of the lepton in the process has also been given

$$\begin{aligned} A(s) &= -3\left(\frac{1-4t^2}{s}\right)^{1/2} E(s)/D(s) \\ E(s) &= \text{Re}(C_8^{eff} C_9^*)s + 2\text{Re}(C_7 C_9^*) + \text{Re}(C_8^{eff} C_{Q_1}^*)t + 2\text{Re}(C_7 C_{Q_1}^*)t \end{aligned} \quad (9)$$

In eqs.(8) and (9) the mass of strange quark has been neglected.

4 Numerical analysis

The constraints on the $SU(5) \times U(1)$ model from $b \rightarrow s\gamma$ have been in detail studied in the ref.[14]. It is shown that there is an upper bound on $\tan\beta$: $\tan\beta \leq 25$ for $\mu > 0$ and there is no analogous bound for $\mu < 0$. The reason is that when $\mu > 0$, the supersymmetric contributions of chargino loops have the same sign with contributions of w and charged higgs loops and interfere constructively, therefore $\tan\beta$ is greatly constrained and the large values of it are not favored; when $\mu < 0$, contributions of SUSY have a opposite sign and interfere destructively, therefore the constraints are relaxed. Because we are interested in the region of large $\tan\beta$, we only consider the case of $\mu < 0$ in the letter.

We analyze the effects of the supersymmetric flipped $SU(5)$ model to the experimental observables — the invariant mass distribution and backward-forward asymmetry. It is obvious from eq.(3) that C_{Q_i} ($i=1, 2$) can reach a value of order one only when $\tan\beta$ is large enough (say, ≥ 20) due to the smallness of $m_b m_\tau / m_h^2$ ($h = h^0, A^0$). For larger values of $\tan\beta$, the contributions of exchanging NHBs dominate, which is shown in fig.2. From the fig.2 one can see the deviations from SM are quite substantial when $\tan\beta=25(30)$ in moduli(dilaton) scenario. The enhancement of the invariant mass distributions is about 200% in both scenarios. The backward-forward asymmetry is greatly modified in both scenarios. The predictions without including the contributions of exchanging NHBs are also shown in fig.2 in order to compare. It is obvious from fig.2 that the invariant mass distribution without including the contributions of exchanging NHBs is almost equal to that in SM because for the values of $\tan\beta$ and $m_{1/2}$ indicated in the fig.2 contributions of SUSY without including those of exchanging NHBs almost do not change the values of C_i ($i=7, 8, 9$) in SM, while for some other values of $\tan\beta$ and $m_{1/2}$, the contributions can change the sign of C_7 and consequently enhance the invariant mass distribution by about 50% compared to SM, similar to the conclusion made in the reference[6].

5 Conclusion

In this paper, we have studied the rare B decay process $B \rightarrow X_s \tau^+ \tau^-$ in the flipped $SU(5)$ model. In particular, the contributions of exchanging neutral higgs bosons is intensively analyzed. It is found that in the regions where $m_{1/2}$ is moderate and $\tan\beta$ is large, the

branching ratio of $B \rightarrow X_s \tau^+ \tau^-$ is enhanced by about 200% compared to the SM and the forward-backward asymmetry is also significantly different from the SM. Therefore, the process $B \rightarrow X_s \tau^+ \tau^-$ is of a good probe to investigate the model and search for new physics.

Acknowledgement

One of the authors (Q.S. Yan) would like to thank W. Liao, H.G. Yan and G.X. Ju for their generous helps and useful discussions. This work was supported in part by the National Natural Science Foundation of China and partly supported by Center of Chinese Advanced Science and Technology(CCAST).

References

- [1] M.F. Sohnius, Phys. Rep. 128 (1985) 39.
- [2] A.B. Lahanas and D.V. Nanopoulos, Phys. Rep. 145 (1987) 1.
- [3] J. Wess and J. Bagger, Supersymmetry and Supergravity, (Princeton Univ, Press 1991).
- [4] G. Jungman, M. Kamionkowski and K. Griest, Phys. Rep. 267 (1996) 195.
- [5] J.L. Lopez and D.V. Nanopoulos, hep-ph/9511266; J.L. Lopez, hep-ph/9601208.
- [6] T. Goto, Y. Okada, Y. Shimizu and M. Tanaka, Phys. Rev. D 55 (1997) 4273.
- [7] ALEPH Collaboration, D. Buskulic et al., Phys. Lett. B 373 (1996) 246; DELPHI Collaboration, P. Abreu et al., ibid. 382 (1996) 323; L3 Collaboration, M. Acciarri et al., ibid. 377 (1996) 289; OPAL Collaboration, G. Alexander et al., ibid, 377 (1996) 181.
- [8] CDF Collaboration, F. Abe et al., Phys. Rev. Lett. 75 (1995) 613; 69 (1992) 3439; D0 Collaboration, S. Abachi et al., ibid. 75, 618(1995).
- [9] Particle Data Group, R.M. Barnett et al., Phys. Rev. D 54 (1996) 1.
- [10] J.P. Derendinger and C.A. Savoy, Nucl. Phys. B 237 (1984) 307; J.A. Casas, A. Lleyda, and C. Munoz, ibid. B 471 (1996) 3; A. Kusenko, P. Lanacker, and G. Segre, Phys. Rev. D 54 (1996) 5824.
- [11] H.E. Habber and G.L. Kane, Phys. Rep. 117 (1985) 75; X. Tata, (304–378) The Standard Model and Beyond (The 9th Symposium on Theoretical Particles, edited by Jihn E. Kim).

- [12] CLEO Collaboration, M.S. Alam et al., Phys. Rev. Lett. 74 (1995) 2885.
- [13] J.L. Lopes, D.V. Nanopoulos and A. Zichichi, CERN-TH 6926/93, CTP-TAMU-33/93, ACT-12/93, Revised; I. Antoniadis, J. Ellis, J. Hagelin and D.V. Nanopoulos, Phys. Lett. B 194 (1987) 231; J. Ellis, J. Hagelin S. Kelley and D.V. Nanopoulos, Nucl. Phys. B 311 (1989) 1.
- [14] J.L. Lopez, D.V. Nanopoulos, X. Wang and A. Zichichi, Phys. Rev. D 51(1995) 147.
- [15] R. Barbieri and G.F. Giudice Phys Lett B 309(1993)86.
- [16] M.A. Diaz, Phys. Lett. B 322 (1994) 591; T. Goto and Y. Okada, Prog. of Theor. Phys. 94 (1995) 407; P. Nath and R. Amorvitt, CERN-TH 7214/94, NUB-TH 3093/94, CTP-TAMU-32/94; R. Garisto and J.N. Ng Phys. Lett. B 315 (1993) 372; J.L. Hewett, SLAC-PUB-6521 May 1994 T/E; F.M. Borzumati, M. Olechowski and S. Pokorski, CERN-TH 7515/94, TUM-73-83/94.
- [17] S. Bertolini, F. borzumati, A. Masieso and G. Ridolfi, Nucl. Phys. B 353 (1991) 591; P. Cho, M. Misiak and D. Wlyer, Phys. Rev. D 54 (1996) 3329.
- [18] J.L. Hewett, Phys. Rev. D 53 (1996) 4964; J.L. Hewett and J.D. Wells Phys. Rev. D 55 (1997) 5549.
- [19] Y.B. Dai, C.S. Huang and H.W. Huang, Phys. Lett. B 390 (1997) 257.
- [20] G.K. Leontaris, Phys. Lett. B 281 (1992) 54.
- [21] S. Kelley, J.L. Lopes and D.V. Nanopoulos, Phys. Lett. B 278 (1992) 140.
- [22] K.R. Dienes, Phys. Rep. 287 (1997) 447.
- [23] V. Kaplunovsky and J. Louis, Phys. Lett. B 306 (1993) 269.
- [24] K. Inoue et al., Prog. Theor. Phys. 68 (1982) 927; A. Bouquet, J. Kaplan and C.A. Savoy, Nucl. Phys. B 262 (1985) 299.
- [25] I.I. Bigi, M. Shifman, N.G. Vraltsev and A.I. Vainstein, Phys. Rev. Lett. 71 (1993) 496; B. Blok, L. Kozrakh, M. Shifman and A.I. Vainstein, Phys. Rev. D 49 (1994) 3356; A.V. Manohar and M.B. Wise, Phys. Rev. D 49 (1994) 1310; S. Balk, T.G. Körner, D. Pirjol and K. Schilcher, Z. Phys. C 64 (1994) 37; A.F. Falk, Z. Ligeti, M. Neubert and Y. Nir, Phys. Lett. B 326 (1994) 145.
- [26] J. Rosiek, Phys. Rev. D 41 (1990) 3464; J.F. Gunion, M.J. Haber, G. Kane and S. Dawson, The Higgs hunter's guide; J.F. Gunion and M.J. Haber, Nucl. Phys. B 272 (1986) 1.
- [27] B. Grinstein, M.J. Savage and M.B. Wise, Nucl. Phys. B 319 (1989) 271.

Figure Captions

Fig.1 The additional Feynman diagrams corresponding to exchanging NHBs in SUSY model, where wavy lines represent the charged bosons: $W^\pm$, $H^\pm$, $\tilde{t}_i$, internal solid lines t quark or charginos, and dashed lines NHBs H^0 , h^0 , A^0 .

Fig.2 $d\Gamma/ds$ and $A(s)$ for the case $\mu < 0$, a) $\tan\beta=25$ and $m_{1/2}=300$ GeV in the moduli scenario and b) $\tan\beta=30$ and $m_{1/2}=350$ GeV in the dilaton scenario. The solid, dashed and dotted lines represent the predictions of the flipped SU(5), the flipped SU(5) without contributions of NHBs and SM respectively.











Fig.1



(a) moduli-scenario, $\tan\beta=25$, $m_{1/2}=300\text{GeV}$



(b) dilaton-scenario, $\tan\beta=30$, $m_{1/2}=350\text{GeV}$

Fig.2: $b \rightarrow s \tau^+ \tau^-$